
REFERENCES

REFERENCES

A

- Abe F., Kato C. & Horikoshi K. (1999). Pressure-regulated metabolism in microorganisms. *Extremophiles in Deep-Sea Environments*, **7**: 447-453.
- Abe F. & Horikoshi K. (2001). The biotechnological potential of piezophiles. *Trends in Biotechnology*, **19**: 102-108.
- Allredge A.L. (1979). The chemical composition of macroscopic aggregates in two neretic seas. *Limnology and Oceanography*, **24**: 855-866.
- Allredge A.L., Gotschalk C., Passow U. & Riebesell U. (1995). Mass aggregation of diatom blooms: Insights from a mesocosm study. *Deep Sea Research II*, **42**: 9-27.
- Ammerman J.W. & Glover W.B. (2000). Continuous underway measurement of microbial ectoenzyme activities in aquatic ecosystems. *Marine Ecology Progress Series*, **201**: 1-12.
- Amon R.M.W. & Benner R. (1996). Bacterial utilisation of different size classes of dissolved organic matter. *Limnology and Oceanography*, **41**: 41-51.
- Amon R.M.W., Fitznar H.-P. & Benner R. (2001). Linkages among the bioreactivity, chemical composition, and diagenetic state of marine dissolved organic matter. *Limnology and Oceanography*, **46**: 287-297.
- Anderson T.R. & Williams P.J. (1999). A one dimensional model of dissolved organic carbon cycling in the water column incorporating combined biological-photochemical decomposition. *Global Biogeochemical Cycles*, **13**: 337-349.
- Avril B.U. (2002). DOC dynamics in the northwestern Mediterranean Sea (DYFAMED site). *Deep Sea Research II*, **49**: 2163-2182.
- Azam F. & Ammerman J.W. (1983). Cycling of organic matter by bacterioplankton in pelagic marine ecosystems: microenvironmental considerations. In *Flows of energy and materials in marine ecosystems. Theory and practice*. M.J.R Fasham (Ed), Plenum Press, New York, p 345-360.
- Azam F., Fenchel T., Field J.G., Gray J.S., Meyer-Reil L.A. & Thingstad F. (1983). The ecological role of water-column microbes in the sea. *Marine Ecology Progress Series*, **10**: 257-263.
- Azam F. & Long R.A. (2001). Sea snow microcosms. *Nature*, **414**: 495-498.

B

- Bartlett D., Wright M., Yayanos A.A. & Silverman M. (1989). Isolation of a gene regulated by hydrostatic pressure in a deep-sea bacterium. *Nature*, **342**: 572-574.
- Bartlett D.H. (1999). Microbial Adaptations to the Psychrosphere/Piezosphere. *Journal of Molecular Microbiology and Biotechnology*, **1**: 93-100.
- Bauer J.E., Williams P.M. & Druffel E.R.M. (1992). ¹⁴C activity of dissolved organic carbon fractions in the north-central Pacific and Sargasso Sea. *Nature*, **357**: 667-670.

- Bélangier C., Desrosiers B. & Lee K. (1997). Microbial extracellular enzyme activity in marine sediments: extreme pH to terminate reaction and sample storage. *Aquatic Microbial Ecology*, **13**: 187-196.
- Benner R., Pakulski J.D., McCarthy M., Hedges J.I. & Hatcher P.G. (1992). Bulk chemical characteristics of dissolved organic matter in the Ocean. *Science*, **25**: 1561-1564.
- Benner R. (2002). Chemical composition and reactivity. In *Biogeochemistry of Marine Dissolved Organic Matter*, D.A. Hansell & C.A. Carlson (Eds), Academic Press, pp. 59-90.
- Berges J.A. & Falkowski P.G. (1998). Physiological stress and cell death in marine phytoplankton: induction of proteases in responses to nitrogen or light limitation. *Limnology and Oceanography*, **43**: 129-135.
- Béthoux J.P., Prieur L. & Bong J.H. (1988). Le courant Ligure au large de Nice. In *Océanographie pélagique méditerranéenne*. J. H. Minas & P. Nival (eds), *Oceanologica Acta*, N° SP 9: 59-67.
- Béthoux J.P. (1989). Oxygen consumption, new production, vertical advection and environmental evolution in the Mediterranean Sea. *Deep-sea Research*, **36**: 769-781.
- Bianchi A. & Garcin J. (1993). In stratified waters the metabolic rate of deep-sea bacteria decreases with decompression. *Deep-Sea Research I*, **40**: 1703-1710.
- Bianchi A. & Garcin J. (1994). Bacterial response to hydrostatic pressure in seawater samples collected in mixed-water and stratified-water conditions. *Marine Ecology Progress Series*, **111**: 137-141.
- Bianchi A., Garcin J., Gorsky G., Poulicek M. & Tholosan O. (1999a). Stimulation du potentiel de dégradation dans les eaux marines profondes par les bactéries barotolérantes colonisant les fèces du plancton migrateur. *Comptes Rendus de l'Académie des Sciences de Paris*, **322**: 1113-1120.
- Bianchi A., Garcin J. & Tholosan O. (1999b). A high-pressure serial sampler to measure microbial activity in the deep sea. *Deep-sea Research I*, **46**: 2129-2142.
- Bianchi A., Tamburini C. & Garcin J. (soumis-a). Microbial activity in deep-sea water: a review of data obtained under *in situ* pressure conditions. *Marine Ecology Progress Series*.
- Bianchi A., Calafat A., De Wit R., Garcin J., Tholosan O., Cacho I., Fabrès J., Grout H., Masqué P., Sanchez-Cabeza J.-A. & Sempéré R. (soumis-b). Microbial activities at the deep water-sediment boundary layer in two high-productivity systems of the Western Mediterranean: The Almeria-Oran Front and the Malaga Upwelling. *Oceanologica Acta*.
- Biddanda B. & Benner R. (1997). Major contribution from mesopelagic plankton to heterotrophic metabolism in the upper ocean. *Deep-Sea Research I*, **47**: 2069-2085.
- Bidle K.D. & Azam F. (1999). Accelerated dissolution of diatom silica by marine bacterial assemblages. *Nature*, **397**: 508-512.
- Bidle K.D. & Azam F. (2001). Bacterial control of silicon regeneration from diatom detritus: significance of bacterial ectohydrolases and species identity. *Limnology and Oceanography*, **46**: 1606-1623.
- Boetius A. & Lochte K. (1994). Regulation of microbial enzymatic degradation of organic matter in deep-sea sediments. *Marine Ecology Progress Series*, **104**: 299-307.

- Borch N.H. & Kirchman D.L. (1997). Concentration and composition of dissolved combined neutral sugars (polysaccharides) in seawater determined by HPLC-PAD. *Marine Chemistry*, **57**: 85-95.
- Brzezinski M.A. & Nelson D.M. (1986). A solvent extraction method for the colorimetric determination of nanomolar concentrations of silicic acid in seawater. *Marine Chemistry*, **19**: 139-151.
- Brooks S.P.J. (1992). A simple computer program with statistical tests for the analysis of enzyme kinetics. *Biotechniques*, **13**: 906-911.
- Buscail R., Pocklington R., Daumas R. & Guidi L. (1990). Fluxes and budget of organic matter in the boundary layer over the northwestern Mediterranean margin. *Continental Shelf Research*, **10**: 1089-1122.
- Buscail R. & Germain C. (1997). Present-day organic matter sedimentation on the NW Mediterranean margin: Importance of off-shelf export. *Limnology and Oceanography*, **42**: 217-229.

C

- Cahet G., Daumas R. & Sibuet M. (1990). In situ experimentation at the water/sediment interface in the deep sea: 2. Biotransformation of dissolved organic substrates by microbial communities at 2000m depth in the Bay of Biscay. *Progress in Oceanography*, **24**: 169-178.
- Cahet G., Gadel F., Martin R. & Bensoussan M. (1981). Remarques sur la réponse de sédiments marins profonds à l'addition de molécules organiques marquées avec mise sous pression. In *Géochimie organique des sédiments marins profonds, Orgon IV. Golfe D'Aden, Mer d'Oman*. R. Pelet (ed.), Editions du CNRS. Paris, pp 61-70
- Carlson C.A., Ducklow H.W. & Michaels A.F. (1994). Annual flux of dissolved organic carbon from euphotic zone in the northwestern Sargasso Sea. *Nature*, **371**: 405-408.
- Carlson C.A. (2002). Production and removal process. In *Biogeochemistry of Marine Dissolved Organic Matter*, D.A. Hansell & C.A. Carlson (Eds), Academic Press: pp. 91-151.
- Caron D.A., Dam H.G., Kremer P., Lessard E.J., Madin L.P., Malone T.C., Napp J.M., Peele E.R., Roman M.R. & Youngbluth M.J. (1995). The contribution of microorganisms to particulate carbon and nitrogen in surface waters of the Sargasso Sea near Bermuda. *Deep-Sea Research I*, **42**: 943-972.
- Cauwet G., Miller A., Brasse S., Fengler G., Mantoura R.F.C. & Spitzky A. (1997). Dissolved and particulate organic carbon in the western Mediterranean Sea. *Deep-Sea Research II*, **44**: 769-779.
- Cho B.C. & Azam F. (1988). Major role of bacteria in biogeochemical fluxes in the ocean's interior. *Nature*, **332**: 441-443.
- Christian J.R. & Karl D.M. (1995). Measuring bacterial ectoenzyme activities in marine waters using mercuric chloride as a preservative and a control. *Marine Ecology Progress Series*, **123**: 217-224.
- Chrost R.J. (1991). Environmental control of the synthesis and activity of aquatic microbial ectoenzymes. In *Microbial enzymes in aquatic environments*, R.J. Chrost (Ed), Springer-Verlag, pp. 29-59.

- Chrost R.J. & Velimirov B. (1991). Measurement of enzyme kinetics in water samples: Effect of freezing and soluble stabilizer. *Marine Ecology Progress Series*, **70**: 93-100.
- Coolen M.J. & Overmann J. (2000). Functional exoenzymes as indicators of metabolically active bacteria in 124,000-year-Old sapropel layers of the eastern Mediterranean Sea. *Applied and Environmental Microbiology*, **66**: 2589-2598.
- Corliss J.B., Dymond J., Gordon L.I., Edmond J.L., Von Erzen R.P., Ballard R.D., Green K., Williams D., Bainbridge A., Crane J., Van Andel T.H. (1979). Submarine thermal springs on the Galapagos Rift. *Science*, **203**: 1073-1083.
- Cowie G.L. & Hedges J.I. (1994). Biochemical indicators of diagenetic alteration in natural organic matter mixtures. *Nature*, **369**: 304-307.

D

- De Bovée F., Guidi L.D. & Soyer J. (1990). Quantitative distribution of deep-sea meiobenthos in the northwestern Mediterranean (Gulf of Lions). *Continental Shelf Research*, **10**: 1123-1145.
- Delmas D. & Garet M.J. (1995). SDS-preservation for deferred measurement of exoproteolytic kinetics in marine samples. *Journal of Microbiological Methods*, **22**: 243-248.
- De Lange G.J. & ten Haven H.L. (1983). Recent sapropel formation in the eastern Mediterranean. *Nature*, **305**: 797-798.
- De Lange G.J., Middelburg J.J., Van der Weijden C.H., Catalano G., Luther G.W., Hydes D.J., Woittiez J.R.W. & Klinkhammer G.P. (1990). Composition of anoxic hypersaline brines in the Tyro and Bannock Basins, Eastern Mediterranean. *Marine Chemistry*, **31**: 63-88.
- DeLong E.F. & Yayanos A.A. (1985). Adaptation of the membrane lipids of a deep-sea bacterium to changes in hydrostatic pressure. *Science*, **228**: 1101-1113.
- DeLong E.F., Franks D. & Yayanos A. (1997). Evolutionary relationships of cultivated psychrophilic and barophilic deep-sea bacteria. *Applied and Environmental Microbiology*, **63**: 2105-2108.
- DeLong E.F., Taylor L.T., Marsh T.L. & Preston C.M. (1999). Visualization and enumeration of marine planktonic archaea and bacteria by using polyribonucleotide probes and fluorescent *in situ* hybridization. *Applied and Environmental Microbiology*, **65**: 5554-5563.
- Deming J.W., Tabor P.S. & Colwell R.R. (1980). Deep ocean microbiology. In *Advanced concepts in Ocean measurements for marine biology*. F. Diemer, J. Vernberg & D. Mirkes (eds), University of South Carolina Press, Columbia St, pp 285-305.
- Deming J.W., Hada H., Colwell R.R., Luehrsen K.R. & Fox G.E. (1984). The ribonucleotide sequence of 5s rRNA from two strains of deep-sea barophilic bacteria. *Journal of Genetic Microbiology*, **130**: 1911-1920.
- Deming J.W. (1985). Bacterial growth in deep-sea sediment trap and boxcore samples. *Marine Ecology Progress Series*, **25**: 305-312.
- Deming J.W., Somers L.K., Straube W.L., Swartz D.G. & MacDonell M.T. (1988). Isolation of an obligately barophilic bacterium and description of a new genus, *Colwellia* gen. nov. *Systematic Applied Microbiology*, **10**: 152-160.

- Deming J.W. (1993). ^{14}C tracer method for measuring microbial activity in deep-sea sediments. In *Handbooks of methods in aquatic microbial ecology*, P.F. Kemp, B.F. Sherr, E.B. Sherr & J.J. Cole (Eds), Lewis Publishers, Boca Raton, Ann Arbor, London, Tokyo, pp 405-414.
- Druffel E.R.M., Williams P.M., Bauer J.E. & Ertel J.R. (1992). Cycling of dissolved and particulate organic matter in the ocean. *Journal of Geophysical Research*, **97**: 15639-15659.
- Ducklow H. (1983). Production and fate of bacteria in the oceans. *BioScience*, **33**: 494-501.
- Ducklow H.W. & Carlson C.A. (1992). Oceanic Bacterial Production. In *Advances in Microbial Ecology*, K.C. Marshall (Ed), Plenum Press, pp. 113-181.
- Dugdale R.C. & Wilkerson F.P. (1998). Silicate regulation of new production in the equatorial Pacific upwelling. *Nature*, **391**: 270-273.

E, F

- Eder W., Jahnke L.L., Schmidt M. & Huber R. (2001). Microbial diversity of the brine-seawater interface of the Kebrat Deep, Red Sea, studied via 16S rRNA gene sequences and cultivation methods. *Applied and Environmental Microbiology*, **67**: 3077-3085.
- Feller G., Narinx E., Arpigny J.L., Aittaleb M., Baise E., Genicot S. & Gerday C. (1996). Enzymes from psychrophilic organisms. *FEMS Microbiology Reviews*, **18**: 189-202.
- Fry J.C. (1990). Direct methods and biomass estimation, *Methods in Microbiology - Volume 22. Techniques in Microbial Ecology*, Lewis Publishers, pp. 41-85.
- Fuhrman J. & Azam F. (1980). Bacterioplankton secondary production estimates for coastal waters of British Columbia, Antarctica, and California. *Applied and Environmental Microbiology*, **39**: 1085-1095.
- Fuhrman J.A. & Davis A.A. (1997). Widespread *Archaea* and novel *Bacteria* from the deep sea as shown by 16S rRNA gene sequences. *Marine Ecology Progress Series*, **150**: 275-285.
- Fukuda R., Ogawa H., Nagata T. & Koike I. (1998). Direct determination of carbon and nitrogen contents of natural bacterial assemblages in marine environments. *Applied and Environmental Microbiology*, **64**: 3352-3358.

G

- Gehlen M., Rabouille C., Ezat U. & Guidi-Guilvard L.D. (1997). Drastic changes in deep-sea sediment porewater composition induced by episodic input of organic matter. *Limnology and Oceanography*, **42**: 980-986.
- Gérino M. (1992). Etude expérimentale de la bioturbation in milieu littoral et profond : Quantification des structures de bioturbation et modélisation du remaniement biologique du sédiment., Université Aix-Marseille II: 208 pp.
- Getliff J.M., Fry J.C., Cragg B.A. & Parkes R.J. (1992). The potential for bacterial growth in deep sediment layers of the Japan Sea, Hole 798B - Leg 128. *Proceeding of the Ocean Drilling Program, Scientific Results*, **127-128**: 755-760.
- Giovannoni S.J., Britschgi T.B., Moyer C.L. & Field K.G. (1990a). Genetic diversity in Sargasso Sea bacterioplankton. *Nature*, **345**: 60-63.

Giovannoni S.J., DeLong E.F., Schmidt T.M. & Pace N.R. (1990b). Tangential flow filtration and preliminary phylogenetic analysis of marine picoplankton. *Applied and Environmental Microbiology*, **56**: 2572-2575.

Giovannoni S. & Rappé M. (2000). Evolution, diversity, and molecular ecology of marine prokaryotes. In *Microbial Ecology of the Oceans*. D. L. Kirchman (ed.), Wiley-Liss, New-York, pp 47-84.

Giuliano L., De Domenico M., De Domenico E., Hofle M.G. & Yakimov M.M. (1999). Identification of Culturable Oligotrophic Bacteria within Naturally Occurring Bacterioplankton Communities of the Ligurian Sea by 16S rRNA Sequencing and Probing. *Microbial Ecology*, **37**: 77-85.

Gray N.D. & Head I.M. (2001). Linking genetic identity and function in communities of uncultured bacteria. *Environmental Microbiology*, **3**: 481-492.

Guillard R.R.L. (1975). Culture of phytoplankton for feeding marine invertebrate animals. In *Culture of marine invertebrate animals*. M. L. Smith & M. H. Chanley (eds), Plenum Press, New York: pp 29-60.

H

Hagström A., Larsson U., Horstedt P. & Normark S. (1979). Frequency of dividing cells, a new approach to the determination of bacterial growth rate in aquatic environments. *Applied and Environmental Microbiology*, **37**: 805-812.

Hansell D.A. & Carlson C.A. (1998). Deep ocean gradients in dissolved organic carbon concentrations. *Nature*, **395**: 263-266.

Hedges J.I., Eglinton G., Hatcher P.G., Kirchman D.L., Arnosti C., Derenne S., Evershed R.P., Kögel-Knabner I., de Leeuw J.W., Littke R., Michaelis W. & Rullkötter J. (2000). The molecularly-uncharacterized component of nonliving organic matter in natural environments. *Organic Geochemistry*, **31**: 945-958.

Hedges J.I., Baldock J.A., Gelinas Y., Lee C., Peterson M. & Wakeham S.G. (2001). Evidence for non-selective preservation of organic matter in sinking marine particles. *Nature*, **409**: 801-804.

Hedges J.I. (2002). Why dissolved organics matter. In *Biogeochemistry of Marine Dissolved Organic Matter*, D.A. Hansell & C.A. Carlson (Eds), Academic Press, pp. 1-33.

Helmke E. & Weyland H. (1991). Effect of temperature on extracellular enzymes occurring in permanently cold marine environments. *Kieler Meeresforschungen, Sonderheft*, **8**: 198-204.

Henneke E. & de Lange G.J. (1990). The distribution of DOC and POC in the water column and brines of the Tyro and Bannock basins. *Marine Chemistry*, **31**: 113-122.

Hoppe H.-G. (1983). Significance of exoenzymatic activities in the ecology of brackish water : measurements by means of methylumbelliferyl-substrates. *Marine Ecology Progress Series*, **11**: 299-308.

Hoppe H.-G. (1993). Use of fluorogenic model substrates for extracellular enzyme activity measurement of bacteria. In *Handbooks of methods in aquatic microbial ecology*, P.F. Kemp, B.F. Sherr, E.B. Sherr & J.J. Cole (Eds), Lewis Publishers, pp. 423-431.

Hoppe H.-G., Ducklow H. & Karrasch B. (1993). Evidence for dependency of bacterial growth on enzymatic hydrolysis of particulate organic matter in the mesopelagic ocean. *Marine Ecology Progress Series*, **93**: 277-283.

Hoppe H.-G. & Ullrich S. (1999). Profiles of ectoenzymes in the Indian Ocean : phenomena of phosphatase activity in the mesopelagic zone. *Aquatic Microbial Ecology*, **19**: 139-148.

Huberton U., Lara R.J. & Kattner G. (1995). Refractory organic compounds in polar waters: relationship between humic substances and amino acids in the Arctic and Antarctic. *Journal of Marine Research*, **53**: 151-166.

J

Jannasch H.W. (1973). Deep-sea microorganisms : *in situ* response to nutrient enrichment. *Science*, **180**: 641-643.

Jannasch H.W. & Taylor C.D. (1984). Deep-sea microbiology. *Annual Review of Microbiology*, **38**: 487-514.

Jansson M., Olsson H. & Petterson K. (1988). Phosphatase: origin, characteristics and function in lakes. *Hydrobiologia*, **170**: 157-175

Jorgensen N.O.G., Kroer N., Coffin R.B. & Hoch M.P. (1999). Relations between bacterial nitrogen metabolism and growth efficiency in an estuarine and an open-water ecosystem. *Aquatic Microbial Ecology*, **18**: 247-261.

K

Karl D.M. & Knauer G.A. (1984). Vertical distribution, transport, and exchange of carbon in the northeast Pacific Ocean: evidence for multiples zones of biological activity. *Deep-Sea Research*, **31**: 221-243.

Karl D.M., Knauer G.A. & Martin J.H. (1988). Downward flux of particulate organic matter in the ocean: a particle decomposition paradox. *Nature*, **332**: 438-440.

Karner M.B., DeLong E.F. & Karl D.M. (2001). Archaeal dominance in the mesopelagic zone of the Pacific Ocean. *Nature*, **409**: 507-510.

Kato C., Suzuki S., Hata S., Ito T. & Horikoshi K. (1995). The properties of a protease activated by high pressure from *Sporosarcina sp.* strain DSK25 isolated from deep-sea sediment. *Jamstecr*, **32**: 7-13.

Kato C., Li L., Nogi Y., Nakamura Y., Tamaoka J. & Horikoshi K. (1998). Extremely barophilic bacteria isolated from the Mariana Trench, Challenger Deep, at a depth of 11,000 meters. *Applied and Environmental Microbiology*, **64**: 1510-1513.

Kato C. (1999). Barophiles (Piezophiles). In K. Horikoshi & K. Tsujii (Eds), *Extremophiles in Deep-Sea Environments*, pp. 91-111.

Kato C. (1999). Molecular analyses of the sediment and isolation of extreme barophiles from the deepest Mariana Trench. In *Extremophiles in Deep-Sea Environments*. K. Horikoshi & K. Tsujii (Eds), pp 27-37.

Kato C. & Qureshi M.H. (1999). Pressure response in deep-sea piezophilic bacteria. *Journal of Molecular Microbiology and Biotechnology*, **1**: 87-92.

- Kato C., Nakasone K., Qureshi M.H. & Horikoshi K. (2000). How do deep-sea microorganisms respond to changes in environmental pressure? *In Environmental Stressors and Gene Responses*. E. b. K. B. S. a. J. Storey (ed.), pp 277-291
- Kato C. & Nogi Y. (2001). Correlation between phylogenetic structure and function: examples from deep-sea *Shewanella*. *FEMS Microbiology Ecology*, **35**: 223-230.
- Keil R.G. & Kirchman D.L. (1991). Dissolved combined amino acids in marine waters as determined by a vapor-phase hydrolysis method. *Marine Chemistry*, **33**: 243-259.
- Keil R.G. & Kirchman D. (1992). Bacterial hydrolysis of protein and methylated protein and its implications for studies of protein degradation in aquatic systems. *Applied and Environmental Microbiology*, **58**: 1374-1375.
- Keil R.G. & Kirchman D.L. (1994). Abiotic transformation of labile protein to refractory protein in sea water. *Marine Chemistry*, **45**: 187-196.
- Keil R.G. & Kirchman D.L. (1999). Utilization of dissolved protein and amino acids in the northern Sargasso Sea. *Aquatic Microbial Ecology*, **18**: 293-300.
- King G.M. & Berman T. (1984). Potential effects of isotopic dilution on apparent respiration in ¹⁴C heterotrophic experiments. *Marine Ecology Progress Series*, **19**: 175-180.
- King G.M. (1986). Characterization of β-glucosidase activity in intertidal marine sediments. *Applied and Environmental Microbiology*, **51**: 373-380.
- Kjørboe T. & Jackson G.A. (2001). Marine snow, organic solute plumes, and optimal chemosensory behavior of bacteria. *Limnology and Oceanography*, **46**: 1309-1318.
- Kirchman D.L., Newell S.Y. & Hodson R.E. (1986). Incorporation versus biosynthesis of leucine : implications for measuring rates of protein synthesis and biomass production by bacteria in marine systems. *Marine Ecology Progress Series*, **32**: 47-59.
- Kirchman D.L., Suzuki Y., Garside C. & Ducklow H.W. (1991). High turnover rates of dissolved organic carbon during spring phytoplankton bloom. *Nature*, **352**: 612-614.
- Kirchman D.L. (1993). Leucine incorporation as a measure of biomass production by heterotrophic bacteria, *Handbooks of methods in aquatic microbial ecology*, P.F. Kemp, B.F. Sherr, E.B. Sherr & J.J. Cole (Eds), Lewis Publishers, pp. 509-512.
- Kirchman D.L. & Ducklow H.W. (1993). Estimating conversion factors for the thymidine and leucine methods for measuring bacterial production. *In Handbook of Methods in Aquatic Microbial Ecology*, P.F. Kemp, B.F. Sherr, E.B. Sherr & J.J. Cole (Eds), Lewis Publishers, pp. 513-517.
- Kirchman D.L. (2002). The ecology of *Cytophaga-Flavobacteria* in aquatic environments. *FEMS Microbiology Ecology*, **39**: 91-100.
- Koike I. & Nagata T. (1997). High potential activity of extracellular alkaline phosphatase in deep waters of the central Pacific. *Deep-Sea Research II*, **44**: 2283-2294.
- Kuznetsova M. & Lee C. (2001). Enhanced extracellular enzymatic peptide hydrolysis in the sea-surface microlayer. *Marine Chemistry*, **73**: 319-332.
- Kyo M., Tuji Y., Usui H. & Itos T. (1991). Collection, isolation and cultivation system for deep-sea microbes study: concept and design. *Ocean*, **1**: 419-423.

L

- Lamy F., Bianchi M., Van Wambeke F., Sempéré R. & Talbot V. (1999). Use of data assimilation techniques to analyze the significance of ectoproteolytic activity measurements performed with the model substrate MCA-Leu. *Marine Ecology Progress Series*, **177**: 27-35.
- Lee S. & Fuhrman J.A. (1987). Relationships between biovolume and biomass of naturally derived marine bacterioplankton. *Applied and Environmental Microbiology*, **53**: 1298-1303.
- Lee C., Wakeham S.G. & Hedges J.I. (2000). Composition and flux of particulate amino acids and chloropigments in equatorial Pacific seawater and sediments. *Deep-Sea Research I*, **47**: 1535-1568.
- Lefèvre D., Denis M., Lambert C.E. & Miquel J.-C. (1996). Is DOC the main source of organic matter remineralization in the ocean water column? *Journal of Marine Systems*, **201**: 281-291.
- Li W.K.W. (1983). Consideration of errors in estimating kinetic parameters based on Michaelis-Menten formalism in microbial ecology. *Limnology and Oceanography*, **28**: 185-190.
- Li L., Kato C., Nogi Y. & Koki H. (1998). Distribution of the pressure-regulated operons in deep-sea bacteria. *FEMS Microbiology Letters*, **159**: 159-166.
- Lochte K. & Turley C.M. (1988). Bacteria and cyanobacteria associated with phytodetritus in the deep sea. *Nature*, **333**: 67-69.
- Lopez-Garcia P., Lopez-Lopez A., Moreira D. & Rodriguez-Valera F. (2001). Diversity of free-living prokaryotes from a deep-sea site at the Antarctic Polar Front. *FEMS Microbiology Ecology*, **36**: 193-202.

M

- McCave I.N. (1975). Vertical flux of particles in the Ocean. *Deep Sea Research*, **22**: 491-502.
- MacDowell M.T. & Colwell R.R. (1985). Phylogeny of the *Vibrionaceae*, and recommendation for two genera, *Listonella* and *Shewanella*. *Systematic Applied Microbiology*, **6**: 171-182.
- Malanotte-Rizzoli P., Manca B.B., d'Alcalà M.R., Theocharis A., Bergamasco A., Bregant D., Budillon G., Civitarese G., Georgopoulos D., Michelato A., Sansone E., Scarazzato P. & Souvermezoglou E. (1997). A synthesis of the Ionian Sea hydrography, circulation and water mass pathways during POEM-Phase I. *Progress in Oceanography*, **39**: 153-204.
- Marty J.-C., Nicolas E., Miquel J.C. & Fowler S.W. (1994). Particulate fluxes of organic compounds and their relationship to zooplankton fecal pellets in the northwestern Mediterranean Sea. *Marine Chemistry*, **46**: 387-405.
- Marty J.-C. & Chiaverini J. (2002). Seasonal and interannual variations in phytoplankton production at DYFAMED time-series station, northwestern Mediterranean Sea. *Deep Sea Research II*, **49**: 2017-2030.
- Marty J.-C., Chiaverini J., Pizay M.-D. & Avril B. (2002). Seasonal and interannual dynamics of nutrients and phytoplankton pigments in the western Mediterranean Sea at the DYFAMED time-series station (1991-1999). *Deep Sea Research II*, **49**: 1965-1985.

- Marxsen J. & Schmidt H.-H. (1993). Extracellular phosphatase activity in sediments of the Breitenbach, a Central European mountain stream. *Hydrobiologia*, **253**: 207-216.
- Massana R., DeLong E.F. & Pedros-Alio C. (2000). A few cosmopolitan phylotypes dominate planktonic archaeal assemblages in widely different oceanic provinces. *Applied and Environmental Microbiology*, **66**: 1777-1787.
- Massana R., Murray A.E., Preston C.M. & DeLong E.F. (1997). Vertical distribution and phylogenetic characterization of marine planktonic Archaea in the Santa Barbara Channel. *Applied and Environmental Microbiology*, **63**: 50-56.
- Mayer L.M. (1989). Extracellular proteolytic enzyme activity in sediments of an intertidal mudflat. *Limnology and Oceanography*, **34**: 973-981.
- McCarthy M., Hedges J. & Benner R. (1996). Major biochemical composition of dissolved high molecular weight organic matter in seawater. *Marine Chemistry*, **55**: 281-297.
- Meyer-Reil L.A. (1986). Measurement of hydrolytic activity and incorporation of dissolved organic substrates by microorganisms in marine sediments. *Marine Ecology Progress Series*, **31**: 143-149.
- Meyer-Reil L.A. (1987). Seasonal and spatial distribution of extracellular enzymatic activities and microbial incorporation of dissolved organic substrates in marine sediments. *Applied and Environmental Microbiology*, **53**: 1748-1755.
- Meyer-Reil L.A. & Köster M. (1992). Microbial life in pelagic sediments: the impact of environmental parameters on enzymatic degradation of organic material. *Marine Ecology Progress Series*, **81**: 65-72.
- Millot C. (1990). The Gulf of Lions' hydrodynamics. *Continental Shelf Research*, **10**: 885-894.
- Miquel J.C., Fowler S.W., La Rosa J. & Buard-Menard P. (1994). Dynamics of the downward flux of particles and carbon in the northwestern Mediterranean Sea. *Deep-Sea Research I*, **41**: 243-261.
- Monaco A., Biscaye A., Soyer J., R. P. & Heussner S. (1990a). Particle fluxes and ecosystem response on a continental margin: the 1985-1988 Mediterranean ECOMARGE experiment. *Continental Shelf Research*, **10**: 809-839.
- Monaco A., Courp T., Heussner S., Carbonne J., Fowler S.W. & Deniaux B. (1990b). Seasonality and composition of particulate fluxes during ECOMARGE-I, western Gulf Lions. *Continental Shelf Research*, **10**: 959-987.
- Mopper K., Zhou X.L., Kieber R.J., Sikorski R.J. & Jones R.D. (1991). Photochemical degradation of dissolved organic carbon and its impact on the organic carbon cycle. *Nature*, **353**: 60-62.
- Moutin T. & Raimbault P. (2002). Primary production, carbon export and nutrients availability in western and eastern Mediterranean Sea in early summer 1996 (MINOS cruise). *Journal of Marine Systems*, **33-34**: 273-288.
- Murray A.E., Preston C.M., Massana R., Taylor L.T., Blakis A., Wu K. & DeLong E.F. (1998). Seasonal and spatial variability of bacterial and archaeal assemblages in the coastal waters near Anvers Island, Antarctica. *Applied and Environmental Microbiology*, **64**: 2585-2595.

- Muyzer G., De Waal E.C. & Uitterlinden A.G. (1993). Profiling of complex microbial population by denaturing gradient gel electrophoresis analysis of polymerase chain reaction-amplified genes coding for 16S rRNA. *Applied and Environmental Microbiology*, **59**: 695-700.
- Muyzer G., Teske A., Wirsén C.O. & Jannasch H.W. (1995). Phylogenetic relationships of *Thiomicrospira* species and their identification in deep-sea hydrothermal vent samples by denaturing gradient gel electrophoresis of 16S rDNA fragments. *Archives of Microbiology*, **164**: 165-172.
- Muyzer G., Brinkhoff U., Nübel C., Sanegoeds H., Schäfer H. & Waver C. (1998). Denaturing gradient gel electrophoresis (DGGE) in microbial ecology. In *Molecular microbial ecology manual*, A. D. L. Akkermans, J. D. van Elsas & F. J. de Bruijn (Eds.), Kluwer Academic Publishers, Dordrecht, The Netherlands, pp 1-27.
- N**
- Nagata T., Fukuda R., Koike I., Kogure K. & Kirchman D.L. (1998). Degradation by bacteria of membrane and soluble protein in seawater. *Aquatic Microbial Ecology*, **14**: 29-37.
- Nagata T., Fukuda H., Fukuda R. & Koike I. (2000). Bacterioplankton distribution and production in deep Pacific waters: Large-scale geographic variations and possible coupling with sinking particle fluxes. *Limnology and Oceanography*, **45**: 426.
- Nelson D.M., Smith W.O.J., Muench R.D., Gordon L.I., Sullivan C.W. & Husby D.M. (1989). Particulate matter and nutrient distributions in the ice-edge zone of the Weddell Sea : relationship to hydrography during late summer. *Deep Sea Research*, **36**: 191-209.
- Nelson D.M., Tréguer P., Brzezinski M.A., Leynaert A. & Quéguiner B. (1995). Production and dissolution of biogenic silica in the ocean: revised global estimates, comparison with regional data and relationship with biogenic sedimentation. *Global Biogeochemical Cycles*, **9**: 359-372.
- Nguyen R.T. & Harvey H.R. (1997). Protein and amino acid cycling during phytoplankton decomposition in oxic and anoxic waters. *Organic Geochemistry*, **27**: 115-128.
- Nikaido H. & Vaara M. (1985). Molecular basis of bacterial outer membrane permeability. *Microbiological Reviews*, **49**: 1-32.
- Nogi Y., Kato C. & Horikoshi K. (1998a). *Moritella japonica* sp. nov., a novel barophilic bacterium isolated from the Japan Trench sediment. *Journal of Genetic Applied Microbiology*, **44**: 289-295.
- Nogi Y., Masui N. & Kato C. (1998b). *Photobacterium profundum* sp. nov., a new, moderately barophilic bacterial species isolated from a deep-sea sediment. *Extremophiles*, **2**: 1-7.
- Nogi Y. & Kato C. (1999). Taxonomic studies of extremely barophilic bacteria isolated from the Mariana Trench and description of *Moritella yayanosii* sp. nov., a new barophilic bacterial isolate. *Extremophiles*, **3**: 71-7.
- Novitsky J.A. (1983). Microbial activity at the sediment-water interface in Halifax Harbor, Canada. *Applied and Environmental Microbiology*, **45**: 1761-1766.

O, P

- Ogawa H., Amagai Y., Koike I., Kaiser K. & Benner R. (2001). Production of refractory dissolved organic matter by bacteria. *Science*, **292**: 917-920.
- Packard T.T. (1979). Half saturation constants for nitrate reductase and nitrate translocation in marine phytoplankton. *Deep-Sea Research*, **26**: 321-326.
- Pantoja S., Lee C. & Marecek J.F. (1997). Hydrolysis of peptides in seawater and sediment. *Marine Chemistry*, **57**: 25-40.
- Pantoja S. & Lee C. (1999). Peptide decomposition by extracellular hydrolysis in coastal seawater and salt marsh sediment. *Marine Chemistry*, **63**: 273-291.
- Picon P. (2000). Diagenèse précoce de la matière organique à l'interface eau-sédiment dans un environnement de marge continentale. Université de Droit, d'Economie et des Sciences d'Aix-Marseille: 221 pp.
- Pomeroy L.R. (1974). The ocean's food webs, a changing paradigm. *BioScience*, **24**: 499-504.
- Porter K.G. & FEIG Y.S. (1980). The use of DAPI for identifying and counting aquatic microflora. *Limnology and Oceanography*, **25**: 943-948.
- Poremba K. (1994a). Impact of pressure on bacterial activity in water columns situated at the european continental margin. *Netherlands Journal of Sea Research*, **33**: 29-35.
- Poremba K. (1994b). Simulated degradation of phytodetritus in deep-sea sediments of the NE Atlantic (47°N, 19°W). *Marine Ecology Progress Series*, **33**: 29-35.
- Poremba K. (1995). Hydrolytic enzymatic activity in deep-sea sediments. *FEMS Microbiology Ecology*, **16**: 213-221.
- Poremba K. & Hoppe H.G. (1995). Spatial variation of benthic microbial production and hydrolytic enzymatic activity down the continental slope of the Celtic Sea. *Marine Ecology Progress Series*, **118**: 237-245.

Q, R

- Qureshi M.H., Kato C. & Horikoshi K. (1998a). Purification of a ccb-type quinol oxidase specifically induced in a deep-sea barophilic bacterium, *Shewanella* sp. strain DB-172F. *Extremophiles*, **2**: 93-99.
- Qureshi M.H., Kato C. & Horikoshi K. (1998b). Purification of two pressure-regulated c-type cytochromes from a deep-sea barophilic bacterium, *Shewanella* sp. strain DB-172F. *FEMS Microbiology Letters*, **161**: 93-99.
- Rath J., Schiller C., & Herndl G.J. (1993). Ecto enzymatic activity and bacterial dynamics along a trophic gradient in the Caribbean Sea. *Marine Ecology Progress Series*, **102**: 89-96.
- Reysenbach A.L., Longnecker K. & Kirshtein J. (2000). Novel bacterial and archaeal lineages from an in situ growth chamber deployed at a Mid-Atlantic Ridge hydrothermal vent. *Applied and Environmental Microbiology*, **66**: 3798-3806.

S

- Sass A.M., Sass H., Coolen M.J.L., Cypionka H. & Overmann J. (2001). Microbial Communities in the Chemocline of a Hypersaline Deep-Sea Basin (Urania Basin, Mediterranean Sea). *Applied and Environmental Microbiology*, **67**: 5392-5402.
- Shanks A.L. & Trent J.D. (1980). Marine snow: sinking rates and potential role in vertical flux. *Deep-Sea Research*, **27A**: 137-143.
- Schmiedl G., de Bovee F., Buscail R., Charriere B., Hemleben C., Medernach L. & Picon P. (2000). Trophic control of benthic foraminiferal abundance and microhabitat in the bathyal Gulf of Lions, western Mediterranean Sea. *Marine Micropaleontology*, **40**: 167-188.
- Sempéré R., Panagiotopoulos C., Lafont R., Marroni B. & Van Wambeke F. (2002). Total organic carbon dynamics in the Aegean Sea. *Journal of Marine Systems*, **33-34**: 355-364.
- Send U., Font J., Krahnemann G., Millot C., Rhein M. & Tintoré J. (1999). Recent advances in observing the physical oceanography of the western Mediterranean Sea. *Progress in Oceanography*, **44**: 37-64.
- Sharp J.H., Benner R., Bennett C.A., Carlson S.E., Fitzwater S.E., Peltzer E.T. & Tupas L.M. (1995). Analysis of organic carbon in seawater: the JGOFS EqPac methods comparison. *Marine Chemistry*, **48**: 91-108.
- Simon M. & Azam F. (1989). Protein content and protein synthesis rates of planktonic marine bacteria. *Marine Ecology Progress Series*, **51**: 201-213.
- Skoog A. & Benner R. (1997). Aldoses in various size fractions of marine organic matter: Implications for carbon cycling. *Limnology and Oceanography*, **42**: 1803-1813.
- Smayda T.J. (1970). The suspension and sinking of phytoplankton in the sea. *Oceanography and Marine Biology Annual Reviews*, **8**: 353-414.
- Smith K.L.J., Carlucci A.F., Jahnke R.A. & Craven D.B. (1987). Organic carbon mineralization in the Santa Catalina Basin: benthic boundary layer metabolism. *Deep-Sea Research*, **36**: 1917-1932.
- Smith D.C., Simon M., Alldredge A.L. & Azam F. (1992). Intense hydrolytic enzyme activity on marine aggregates and implications for rapid particle dissolution. *Nature*, **359**: 139-142.
- Smith D.C., Steward G.F., Long R.A. & Azam F. (1995). Bacterial utilization of carbon fluxes during a diatom bloom in a mesocosm. *Deep Sea Research II*, **42**: 75-97.
- Somville M. & Billen G. (1983). A method for determining exoproteolytic activity in natural waters. *Limnology and Oceanography*, **28**: 190-193.
- Stahl D.A., Lane D.J., Olsen G.J. & Pace N.R. (1984). Analysis of hydrothermal vent associated symbiots by ribosomal RNA sequences. *Science*, **244**: 409-411.
- Stemann L., Gorsky G., Marty J.-C., Picheral M. & Miquel J.-C. (2002). Four-year study of large-particle vertical distribution (0-1000m) in the NW Mediterranean in relation to hydrology, phytoplankton, and vertical flux. *Deep Sea Research II*, **49**: 2143-2162.
- Stepanuskas R., Edling H. & Tranvik L.J. (1999). Differential dissolved organic nitrogen availability and bacterial aminopeptidase activity in limnic and marine waters. *Microbial Ecology*, **38**: 264-272.
- Stewart W.W. (1981). Synthesis of 3,6-disulfonated 4-aminonaphthalimides. *Journal Of American Chemical Society*, **103**: 7614-7620.

- Strickland J.D.H. & Parsons T.R. (1972). A Practical Handbook of Seawater Analysis. *Fish. Res. Bd. Can. Bull.*, **167**: 1-310.
- Sugimura Y. & Suzuki Y. (1988). A high-temperature catalytic oxidation method for the determination of non-volatile dissolved organic carbon in seawater by direct injection of liquid sample. *Marine Chemistry*, **24**: 105-131.
- Suzuki Y., Sugimura Y. & Itoh T. (1985). A catalytic oxidation method for the determination of total nitrogen dissolved in seawater. *Marine Chemistry*, **16**: 83-97.

T

- Takai K. & Horikoshi K. (1999). Genetic diversity of archaea in deep-sea hydrothermal vent environments. *Genetics*, **152**: 1285-1297.
- Talbot V. & Bianchi M. (1997). Bacterial proteolytic activity in sediments of the Subantarctic India Ocean sector. *Deep-Sea Research II*, **44**: 1069-1084.
- Talbot V., Giuliano L., Bruni V. & Bianchi M. (1997). Bacterial abundance, production and ectoproteolytic activity in the Strait of Magellan. *Marine Ecology Progress Series*, **154**: 293-302.
- Tamburini C., Garcin J., Ragot M. & Bianchi A. (2002). Biopolymer hydrolysis and bacterial production under ambient hydrostatic pressure through a 2000 m water column in the NW Mediterranean. *Deep Sea Research II*, **49**: 2109-2123.
- Tamburini C. & Tedetti M. (soumis-a). A simple and highly reproducible to measure the bacterial respiration of ^{14}C -labeled compounds in seawater samples. *Hydrobiologia*.
- Tamburini C., Garcin J. & Bianchi A. (soumis-b). Microbial life in the deep anoxic hypersaline basins of the Eastern Mediterranean. *Nature*.
- Tamburini C., Garcin J. & Bianchi A. (soumis-c). Role of deep-sea bacteria in organic matter mineralization and adaptation to hydrostatic pressure conditions in the NW Mediterranean Sea. *Aquatic Microbial Ecology*.
- Tanaka T. & Rassoulzadegan F. (2002). Full-depth profile (0-2000 m) of bacteria, heterotrophic nanoflagellates and ciliates in the NW Mediterranean Sea: Vertical partitioning of microbial trophic structures. *Deep Sea Research II*, **49**: 2093-2107.
- Tengberg A., De Bovee F., Hall P., Berelson W., Chadwick D., Ciceri G., Crassous P., Devol A., Emerson S., Gage J., Glud R., Graziottini F., Gundersen J., Hammond D., Helder W., Hinga K., Holby O., Jahnke R., Khripounoff A., Lieberman S., Nuppenau V., Pfannkuche O., Reimers C., Rowe G., Sahami A., Sayles F., Schurter M., Smallman D., Wehrli B. & De Wilde P. (1995). Benthic chamber and profiling landers in oceanography - A review of design, technical solutions and functioning. *Progress in Oceanography*, **35**: 253-294.
- Tholosan O. & Bianchi A. (1998). Bacterial distribution and activity at the water-sediment boundary layer on NW Mediterranean continental margin. *Marine Ecology Progress series*, **168**: 273-283.
- Tholosan O. (1999). Activités microbiennes dans les eaux et les sédiments profonds. Rôle de la pression hydrostatique. Université de la Méditerranée, pp. 275.

- Tholosan O., Lamy F., Garcin J., Polychronaki T. & Bianchi A. (1999). Biphasic Extracellular Proteolytic Enzyme Activity in Benthic Water and Sediment in the Northwestern Mediterranean Sea. *Applied and Environmental Microbiology*, **65**: 1619-1626.
- Tréguer P., Kamatani A., Gueneley S. & Quéguiner B. (1989). Kinetics of dissolution of Antarctic diatom frustules and the biogeochemical cycle of silicon in the Southern Ocean. *Polar Biology*, **9**: 397-403.
- Tréguer P., Nelson D.M., Van Bennekom A.J., Demaster D.J., Leynaert A. & Quéguiner B. (1995). The balance of silica in the world ocean: a re-estimate. *Science*, **268**: 375-379.
- Turley C., & Lochte K. (1990). Microbial response to the input of fresh detritus to the deep-sea bed. *Palaeogeography, Palaeoclimatology, Palaeoecology*, **89**: 3-23.
- Turley C.M. (1993). The effect of pressure on leucine and thymidine incorporation by free-living bacteria and by bacteria attached to sinking oceanic particles. *Deep-Sea Research I*, **40**: 2193-2206.
- Turley C.M. & Mackie P.J. (1994). Biogeochemical significance of attached and free-living bacteria and the flux of particles in the NE Atlantic Ocean. *Marine Ecology Progress Series*, **115**: 191-203.
- Turley C.M., Lochte K. & Lampitt R.S. (1995). Transformation of biogenic particles during sedimentation in the northeastern Atlantic. *Phil. Trans. R. Soc. Lond. N*, **348**: 179-189.
- Turley C.M., Lochte K. & Patterson D.J. (1988). A barophilic flagellate isolated from 4500 m in the mid-North Atlantic. *Deep-Sea Research*, **35**: 1079-1092.

U, V

- Unanue M., Ayo B., Slezak D., Herndl G.J. & Ibarri J. (1999). Ecto-enzymatic activity and uptake of monomers in marine bacterioplankton described by a biphasic kinetic model. *Microbial Ecology*, **37**: 36-48.
- Van Wambeke F. (1988). Numération des bactéries planctoniques au moyen de l'analyse d'image couplée à l'épifluorescence. *Annales de l'Institut Pasteur/Microbiologie*, **139**: 261-272.
- Van Wambeke F. (1995). Numération et taille des bactéries planctoniques au moyen de l'analyse d'images couplée à l'épifluorescence. *Océanis*, **21**: 113-124.
- Van Wambeke F., Goutx M., Striby L., Sempéré R. & Vidussi F. (2001). Bacterial dynamics during the transition from spring bloom to oligotrophy in the northwestern Mediterranean Sea: relationships with detritus and dissolved organic matter. *Marine Ecology Progress Series*, **212**: 89-105.
- Vetriani C., Jannash H.W., McGregor B.J., Stahl D.A. & Reysenbach A.-L. (1999). Population structure and phylogenetic characterization of marine benthic Archaea in deep-sea sediments. *Applied and Environment Microbiology*, **65**: 4375-4384.

W, Y, Z

- Wakeham S.G. & Lee C. (1989). Organic geochemistry of particulate organic matter in the ocean: the role of particles in oceanic sedimentary cycles. *Organic Geochemistry*, **14**: 83-96.

- Wakeham S.G., Lee C., Hedges J.I., Hernes P.J. & Peterson M.L. (1997). Molecular indicators of diagenetic status in marine organic matter. *Geochimica et Cosmochimica Acta*, **61**: 5363-5369.
- Weinbauer M.G., & Höfle M.G. (1998). Distribution and life strategies of two bacterial populations in a eutrophic lake. *Applied and Environmental Microbiology*, **64**: 3776-3783.
- Williams P.J.L.B. & Gray R.W. (1970). Heterotrophic utilization of dissolved organic compounds in the sea. II. Observations on the responses of heterotrophic marine populations to abrupt increases in amino acid concentration. *Journal marine biologists Association U.K.*, **50**: 871-881.
- Williams P.M. & Druffel E.R.M. (1987). Radiocarbon in dissolved organic matter in the central North Pacific Ocean. *Nature*, **330**: 246-248.
- Williams B.J.I. (2000). Heterotrophic bacteria and the dynamics of dissolved organic material. In *Microbial Ecology, of the Oceans*, D.L. Kirchman (Ed), Wiley-Liss, pp. 153-200.
- Wright R.T. & Hobbie J.E. (1965). The uptake of organic solutes in lake water. *Limnology and Oceanography*, **10**: 22-28.
- Wright R.T. & Hobbie J.E. (1966). Use of glucose and acetate by bacteria and algae in aquatic ecosystems. *Ecology*, **47**: 447-464.
- Yanagibayashi M., Nogi Y., Li L. & Kato C. (1999). Changes in the microbial community in Japan Trench sediment from a depth of 6292 m during cultivation without decompression. *FEMS Microbiological Letters*, **170**: 271-279.
- Yayanos A.A. (1995). Microbiology to 10,500 Meters in the Deep Sea. *Annual Review of Microbiology*, **49**: 777-805.
- Yayanos A.A., S. D.A. & Van Boxtel R. (1979). Isolation of a deep-sea barophilic bacterium and some of its growth characteristics. *Science*, **205**: 808-809.
- ZoBell C.E. & Johnson F.H. (1949). The influence of hydrostatic pressure on the growth and viability of terrestrial and marine bacteria. *Journal of Bacteriology*, **57**: 179-189.
- ZoBell C.E. (1970). Pressure effects on morphology and life processes of bacteria. In H.M. Zimmerman (Ed), *High pressure effects on cellular processes*, Academic Press, pp. 85-130.
- Zweiffel U.L. & Hagström A. (1995). Total counts of marine bacteria include a large fraction of non-nucleoid-containing bacteria (ghosts). *Applied and Environmental Microbiology*, **61**: 2180-2185.